

# Parameterization of Soil Respiration in GEMTM

Erica McGrath-Spangler

AT 730

Spring 2006

# Outline

- Equation
- Components of the Parameterization
- Parameter Values
- Original Equation
- Parameter  $a'$
- Parameter  $T_{s,ref}$
- Parameter  $c'$
- Temperature Sensitivity
- Soil Water Content Sensitivity
- Look up Table
- Summary

# Equation

- Of the form:

$$F_{\text{css}} = a' (\Theta_{20} - 12) / (40 - 12) e^{c'(T_{s,10} - T_{s,\text{ref}})}$$

- $F_{\text{css}}$  is the soil respiration rate in  $\mu\text{mol m}^{-2}\text{s}^{-1}$
- Function of available carbon, soil moisture, and soil temperature

# Equation

$$F_{\text{css}} = a' (\Theta_{20} - 12) / (40 - 12) e^{c'(T_{s,10} - T_{s,\text{ref}})}$$

- $a'$  is the soil  $\text{CO}_2$  flux at field capacity
- Describes how much carbon is available for decomposition

# Equation

$$F_{\text{css}} = a' \left( \frac{\Theta_{20} - 12}{40 - 12} \right) e^{c'(T_{s,10} - T_{s,\text{ref}})}$$

- $\Theta_{20}$  is the soil-water content in percent at 20 cm depth
- Describes the part of the function dependent upon soil moisture

# Equation

$$F_{\text{css}} = a'(\Theta_{20} - 12)/(40-12)e^{\frac{c'(T_{s,10} - T_{s,\text{ref}})}{10}}$$

- $c'$  is the temperature coefficient and determines the soil respiration's dependence on temperature
- Describes the function's dependence upon temperature

# Equation

$$F_{\text{css}} = a'(\Theta_{20} - 12)/(40-12)e^{c'(T_{s,10} - T_{s,\text{ref}})}$$

- $T_{s,10}$  is the soil temperature in °C at 10cm depth
- $T_{s,\text{ref}}$  is the reference soil temperature at 10cm depth

# Equation

$$F_{\text{css}} = a'(\Theta_{20} - 12)/(40-12)e^{c'(T_{s,10} - T_{s,\text{ref}})}$$

- 12% is the soil water content when soil CO<sub>2</sub> fluxes go to zero - this is just drier than the permanent wilting stage
- 40% is near field capacity - when the prescribed CO<sub>2</sub> fluxes occur

# Components of the parameterization

<u>Input</u>	<u>Output</u>	<u>Parameters</u>
$\Theta_{20}$	$F_{\text{css}}$	$a'$
$T_{s,10}$		$c'$
		$T_{s,\text{ref}}$
		12
		40

# Parameter Values

- Based off of 1000 observations of soil surface CO<sub>2</sub> fluxes in the FIFE area (central Kansas), Norman et al., 1992 set
  - $a' = 12.1$
  - $c' = 0.0365$
  - $T_{s,ref} = 26.0$

# Parameter Values

- However, for these observations,  $T_{s,10}$  varied only between 20°C and 30°C
- Another study by Grammerer, 1989 made observations between 3°C and 30°C which gave different values for  $c'$  and  $T_{s,ref}$
- Changes  $c' = 0.069$ ,  $T_{s,ref} = 25$ , and  $a' = 11$  to best fit the 1000 observations with the new  $c'$  and  $T_{s,ref}$  based on Grammerer's study

# Original Equation

- Forms the equation

$$F_{\text{css}} = 11(\Theta_{20} - 12)/(40-12)e^{0.069(T_{s,10} - 25)}$$


- This form of the equation was used in GEMTM when it was first built
- However, the value of a' was based only on one vegetation type (tall grass) in central Kansas

# Parameter a'

- The parameter a' is the product of the heterotrophic respiration rate  $K_d$  at  $0^\circ\text{C}$  and  $C_s$ , the carbon in the soil and detritus (dead or decaying organic matter)
- At present in GEMTM, a' is based off of 28 different vegetation types with values ranging from 0.4161 to 6


# Parameter a'

- The blue represents the range of values of a' used in GEMTM
- The red is if the value of  $K_d$  and  $C_s$  are each increased by 10%
- The yellow is if the values of  $K_d$  and  $C_s$  are each decreased by 10%



# Parameter a'

- 1 evergreen needleleaf tree
- 2 deciduous needleleaf tree
- 3 deciduous broadleaf tree
- 4 evergreen broadleaf tree
- 5 short grass
- 6 tall grass
- 7 desert
- 8 semi-desert
- 9 tundra
- 10 evergreen shrub
- 11 deciduous shrub
- 12 mixed woodland
- 13 crop/mixed farming
- 14 irrigated crop
- 15 bog or marsh
- 16 evergreen needleleaf forest
- 17 evergreen broadleaf forest
- 18 deciduous needleleaf forest
- 19 deciduous broadleaf forest
- 20 mixed cover
- 21 woodland
- 22 wooded grassland
- 23 closed shrubland
- 24 open shrubland
- 25 grassland
- 26 cropland (corn)
- 27 bare ground
- 28 urban and built up



# Parameter a'

- The default in GEMTM uses vegetation types 1 - 15,22,27, and 28
- Many of the different vegetation types use the same value for a'


- 1 evergreen needleleaf tree
- 2 deciduous needleleaf tree
- 3 deciduous broadleaf tree
- 4 evergreen broadleaf tree
- 5 short grass
- 6 tall grass
- 7 desert
- 8 semi-desert
- 9 tundra
- 10 evergreen shrub
- 11 deciduous shrub
- 12 mixed woodland
- 13 crop/mixed farming
- 14 irrigated crop
- 15 bog or marsh
- 16 evergreen needleleaf forest
- 17 evergreen broadleaf forest
- 18 deciduous needleleaf forest
- 19 deciduous broadleaf forest
- 20 mixed cover
- 21 woodland
- 22 wooded grassland
- 23 closed shrubland
- 24 open shrubland
- 25 grassland
- 26 cropland (corn)
- 27 bare ground
- 28 urban and built up

# Parameter $a'$

- $a'$  is very uncertain and varies according to soil type and vegetation
- Another set of values comes from Raich and Schlesinger, 1992
  - Uses more values from North America, Europe, and Asia


# Parameter a'

- Vegetation type
- 1 tundra
  - 2 boreal forests and woodlands
  - 3 temperate grasslands
  - 4 temperate coniferous forests
  - 5 temperate deciduous forests
  - 6 mediterranean woodlands and heath
  - 7 croplands, fields, etc.
  - 8 desert scrub
  - 9 tropical savannas and grasslands
  - 10 tropical dry forests
  - 11 tropical moist forests
  - 12 northern bogs and mires
  - 13 marshes



# Parameter a'

FIFE




QuickTime™ and a  
TIFF (Uncompressed) decompressor  
are needed to see this picture.

# Parameter $T_{s,ref}$

- Depends on the observations and the soil temperatures of the observations
- Initially taken to be 26°C based on soil temperatures between 20°C and 30°C
- Changed to 25°C based on temperatures between 3°C and 30°C

# Parameter $T_{s,ref}$

- As temperature increases, the dependence on  $T_{s,ref}$  increases
- A 10% decrease in  $T_{s,ref}$  can lead to a 20% increase in respiration due to the exponential




# Parameter $c'$

- The value of  $c'$  also depends on the observations and the soil temperatures of the observations
- Initially,  $c' = 0.0365$  based on temperatures between  $20^{\circ}\text{C}$  and  $30^{\circ}\text{C}$  from Norman et al., 1992
- Changed to  $0.069$  based on observations between  $3^{\circ}\text{C}$  and  $30^{\circ}\text{C}$  by Grammerer, 1989


# Parameter $c'$

- Red colors are air temperatures greater than  $30^{\circ}\text{C}$  on 21 April 2006 at 19:00 UTC
- For these regions the temperature range for which  $c'$  was calculated may not apply depending on how these warm temperatures transfer through the soil




# Parameter $c'$

- Similar issue appears in Africa
- Is  $c'$  valid for these warm regions?



# Parameter $c'$

- For soil temperatures very near the reference temperature, variations in  $c'$  are unimportant
- At temperatures away from the reference temperature, variations in  $c'$  can make a large difference since  $c'$  is in an exponential



# Temperature Sensitivity

- The temperature sensitivity of soil respiration can be affected by:
  1. Physical protection
  2. Chemical protection
  3. Drought
  4. Flooding
  5. Freezing

# Temperature Sensitivity

- Physical protection
  - Organic matter physically protected in the interior of soil aggregates; microorganisms and enzymes have limited access
  - Climate can affect aggregate formation through the action of raindrops and the growth of fungal hyphae

# Temperature Sensitivity

- Chemical Protection
  - Organic matter adsorbed onto mineral surfaces through bonds
  - This process also affected by temperature

# Temperature Sensitivity

- Drought
  - Reduces the thickness of soil water films, inhibiting diffusion of extracellular enzymes and soluble organic-C substrates
  - Determined by climate-driven hydrologic balance

# Temperature Sensitivity


- Flooding
  - Slows oxygen diffusion, allowing only anaerobic decomposition which is generally slower
  - Flooding determined by precipitation and evapotranspiration

# Temperature Sensitivity

- Freezing
  - Diffusion of substrates and extracellular enzymes is slow when the soil water is frozen
  - Melting of permafrost will expose additional organic matter


# Soil Water Content Sensitivity

- Relationship of soil water content to respiration is linear in GEMTM





# Soil Water Content Sensitivity

- In other models, relationship is different
- In dry regions, drought prevents much microbial activity in the soil
- When too much soil water content, microbes drown and cannot decompose soil organic matter and so produces a reduction in soil respiration



# Soil Water Content Sensitivity

- In GEMTM, soil respiration continues to increase as soil water content increases
- Neglects the affect of flooding and limitation to anaerobic respiration only



# Look up Table

- Much of the parameterization is already in the form of a look up table due to the dependence of  $a'$  on biome type
- The model knows what the simulated vegetation type is and goes to an array of values and picks out the  $a'$  that corresponds to that type

# Look up Table

- Could have a look up table for the exponential function
- The rest of the parameterization is linear and not much would be gained by a look up table
- The computer would not need to recompute exponentials every time it runs the program for each temperature
- Accuracy of this method depends very much on temperature and becomes less accurate as temperature increases

# Look up Table


Temperature	$\exp(0.069*(T-25))$
0	0.178173052
1	0.190901059
2	0.204538307
3	0.219149748
4	0.234804976
5	0.251578553
6	0.269550371
7	0.288806028
8	0.309437236
9	0.331542259
10	0.355226381
11	0.380602407
12	0.4077912
13	0.436922258
14	0.468134327
15	0.501576069

Temperature	$\exp(0.069*(T-25))$
16	0.537406762
17	0.575797064
18	0.616929823
19	0.661000951
20	0.708220353
21	0.758812931
22	0.81301965
23	0.871098692
24	0.93332668
25	1
26	1.071436209
27	1.14797555
28	1.229982572
29	1.317847864
30	1.41198992

# Summary

- Soil respiration in GEMTM based on 5 parameters
- Format of current parameterization based on observations in central Kansas with limited soil temperature, soil type, and vegetation type
- Parameters are highly variable depending on temperature, soil type, and vegetation type
- Need different values for different geographical locations
- Soil respiration is much more complex than what is accounted for in parameterizations

# Image of soil CO<sub>2</sub> emissions



# References

- Chen, D.X., and Coughenour, M.B., 1994. GEMTM: a general model for energy and mass transfer of land surfaces and its application at the FIFE sites. *Agricultural and Forest Meteorology*, 68: 145-171
- Davidson, E.A. and Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*, 440: 165-173.
- Norman, J.M., Garcia, R. and Verma, S.B., 1992. Soil surface CO<sub>2</sub> fluxes and carbon budget of a grassland. *J. Geophys. Res.*, 97 (D17): 18845-18853.
- Raich, J.W., Rastetter, E.B., Melillo, J.M., Kicklighter, D.W., Steudler, P.A., Parton, B.J., Grace, A.L., Moore III, B., and Vorosmarty, C.J., 1991. Potential net primary productivity in South America: Application of a global model. *Ecological Applications*, 1(4): 399-429.
- Raich, J.W. and Schlesinger, W.H., 1992. The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus*, 44B: 81-99.

A BIG thanks to Adriana Beltran!!